



SPECIFIC METHODS OF THE GAMMA FUNCTION

Tog'ayev Turdimurod Xurram ug'li

turdimurodtagayev@gmail.com

Xolboev Sanjar Boboxonovich

xolboyevsanjar0911@gmail.com

Jo'rayev Ixhtiyor Baxtiyor o'gli

ixhtiyorjurayev05@gmail.com

Hamidova Sevinch Safarmamat qizi

sevinchhamidova789@gmail.com

Denau Institute of Entrepreneurship and Pedagogy

<https://doi.org/10.5281/zenodo.15576811>

Annotation

This article explores Euler integrals, specifically the gamma and beta functions, and their significance in mathematical analysis, with applications in differential equations, probability theory, and statistics. The gamma function, defined as a continuous extension of the factorial, is analyzed for its key properties, including continuity, infinite differentiability, and the functional relation $\Gamma(z+1)=z\Gamma(z)$. Singular points ($t=0$ and $t=\infty$) and the uniform convergence of the integral are examined using the Weierstrass criterion. Applications of the gamma function in probability distributions, quantum mechanics, and statistical physics are highlighted through examples. The article also includes a practical integral calculation and is supported by references to mathematical literature.

Keywords: Gamma function, Euler integrals, beta function, probability theory, statistics, differential equations.

Euler Integrals.

Euler integrals are among the fundamental mathematical tools that include gamma and beta functions. Both functions represent continuous functions and special processes, and are widely applied in the fields of differential equations, probability theory, and statistics.

Gamma Function

The gamma function is particularly widely used in probability distributions, such as the gamma distribution. This function is a continuous equivalent of the factorial and is employed in numerous statistical models. For instance, in the process of multidimensional statistical modeling, distributions and integral calculations are generated using the gamma function. In physics, the gamma function also serves as an essential tool for models in quantum mechanics and statistical physics. The gamma function is a function that generalizes factorials and is defined as follows:

$$\Gamma(z) = \int_0^{+\infty} t^{z-1} e^{-t} dt, \quad \text{Re}(z) > 0. \quad (1)$$

The points $t=0$ and $t=\infty$ are called singular points. Let's express the integral (1) as the sum of two integrals:

$$\Gamma(z) = \int_0^1 t^{z-1} e^{-t} dt + \int_1^{+\infty} t^{z-1} e^{-t} dt$$

These two integrals are uniformly convergent in any finite interval with respect to the parameter according to the Weierstrass criterion. Indeed, if:

Properties of the gamma function.

. (1) The gamma function is continuous for all values of the argument and has continuous derivatives of all orders. By differentiating the integral (1) under the integral sign,

$$\Gamma' = \int_1^{+\infty} t^{z-1} \ln t e^{-t} dt \text{ ni hosil qilamiz.}$$

$\int_0^1 t^{z-1} \ln t e^{-t} dt$ va $\int_1^{+\infty} t^{z-1} \ln t e^{-t} dt$ Since the integrals are uniformly convergent over an arbitrary interval, it is possible to apply Leibniz's rule:

For the first case, when $t=0$, the majorant function is [missing information], while for the second case, when [missing information], the majorant function is [missing information]. Based on induction, it can be shown that $\Gamma(z)$ is an infinitely differentiable function for $z>0$.

$$\Gamma^{(n)}(z) = \int_1^{+\infty} t^{z-1} (\ln t)^n e^{-t} dt \quad (2)$$

2⁰. Now let us derive the basic functional relation for the Gamma function. Let . As a result of integration by parts, the following

$$\Gamma(z+1) = \int_1^{+\infty} t^z e^{-t} dt = -e^t t^z \Big|$$

$$\Gamma(z+1) = z \Gamma(z) \quad (3)$$

ni hosil qilamiz. (18) formula pasaytirish formulasi deyiladi. Bu formuladan xususiylar uchun

$$\Gamma(n+1) = n!$$

We take the formula. From formula (3), it is possible to determine the behavior of the function $\Gamma(z)$ with respect to z : since $\Gamma(z+1)$ is a continuous function and because it is

$$\Gamma(z) = \frac{\Gamma(z+1)}{z} \sim \frac{\Gamma(1)}{z}.$$

Formula (3) allows for the extension of the $\Gamma(z)$ function, while preserving its properties, even for negative values of z that are not equal to $-1, -2, -3, \dots, -n, \dots$

$$\Gamma(z) = \frac{\Gamma(z+1)}{z} \quad (4)$$

will be. Since $-1 < z < 0$, the definition (4) is correct. Let's consider the character of the function at $y=z+1$. Taking $y=z+1$, we obtain an equivalent function to $y=z+1$. Therefore, instead of (4) in

$$\Gamma(y-1) = \frac{\Gamma(y)}{y-1} \sim -\Gamma(y) \sim -\frac{1}{y} = -\frac{1}{z+1};$$

Let's consider this. Therefore, $z \rightarrow -1 + 0$ da $\Gamma(z) \sim -\frac{1}{z+1}$, mainly $z \rightarrow -n$ da $\Gamma(z) \sim -\frac{(-1)^n}{z+n}$ it can be determined..[1-15]

Example, $\int_0^{+\infty} x^{2n} e^{-x^2} dx$, $n \in \mathbb{N}$, calculate the integral.

Solution: $x \rightarrow \sqrt{t}$, ($t>0$) saying that,

$$\int_0^{+\infty} x^{2n} e^{-x^2} dx = \frac{1}{2} \int_0^{+\infty} t^{n-\frac{1}{2}} e^{-t} dt = \frac{1}{2} \Gamma(n+2) = \frac{(2n+1)!}{2^{n+1}} \pi;$$

we produce..

References

References:

- 1.A.U.Abdurhamidov, H.A.Nasimov, U.M.Nosirov, J.H.Husanov "Algebra va matematik analiz asoslari" 1-2 qism. "O'quvchi".T.2008.
- 2.Nassiet S , Torte D , Rivoallan L , Matematik Analiz. Didier, Paris, 1995. 3. Alimov Sh.A va b. Algebra va analiz asoslari, 10-11. "O'qituvchi", 1996 . 4. <http://www.ziyonet.uz>
- 3.В.В.Вавилов, Н.И.Мельников, С.Н.Олехник, ПасиченкоП.Н. Задачи по математике . Алгебра. Справочной пособие. Москва.Наука.1987г.
- 4.С.Н.Олехник и др. Уравнения и неравенства.Нестандартные методы решения. Учебно - метод. пособие. Москва. 2001.
- 5.Mirzaahmedov M. va boshqalar. Matematikadan olimpiada masalalari. Toshkent. O'qituvchi. 1997.
- 6.Abdurashidov Nuriddin, Tog'ayev Turdimurod, Rustamov Bilol, Eshtemirov Eshtemir "Equation of the Result of Second-Order Surfaces" EXCELLENCIA: INTERNATIONAL MULTI-DISCIPLINARY JOURNAL OF EDUCATION <https://multijournals.org/index.php/excellencia-imje>
- 7.Abdurashidov Nuriddin G.,iyoziddin o.,g.,li ,Tog.,ayev Turdimurod Xurram o.,g.,li,Rustamov Bilol Muxbiddinovich. "Laplas tenglamasining fundamental yechimi" . "So'ngi ilmiy tadqiqotlar nazariyasi" Ilmiy-uslubiy jurnali. 13-iyun 2024- yil 7-jild 6-son (33-37).
- 8.Abdurashidov Nuriddin , Tog'ayev Turdimurod, Eshtemirov Eshtemir, Toshtemirova Sarvara. Laplas teoremasi yordamida 5-tartibli determinantni hisoblash. "O'ZBEKISTONDA FANLARARO INNOVATSIYALAR VA ILMIY TADQIQOTLAR" 20-dekabr 2024-yil 35-son (343-347)
- 9.Салоҳиддинов М. Математик физика тенгламалари. Тошкент-«Ўзбекистон» нашриёти-2002й.-448 б. 2. Салахитдинов МС, Мирсабуров. Нелокальные задачи для уравнений смешанного типа с ...
https://scholar.google.com/citations?view_op=view_citation&hl=en&user=7ZmvhwsAAAAJ&citation_for_view=7ZmvhwsAAAAJ:9yKSN-GCB0IC
- 10.B.M.Rustamov. J.U.To'raxonov. Kasodlik ehtimolligining aniq hisobiga doir misol. Образование наука и инновационные идеи в мире 35(2) 172-175
- 11.Rustamov Bilol, Baltabayeva Saida, Choriyeva Munira,Normo,minova Charos "" Diskret tasodifiy miqdorning sonli xarakteristikalari"."O'zbekistonda fanlararo innovatsiyalar va ilmiy tadqiqotlar " 20-fevral 2025- yil 37-son (292-297).
- 12.B.M.Rustamov, N.G'.Abdurashidov, Sh.Ashirov, A.Saitniyozov. "" International Journal of Education, Social Science & Humanities. Finland Academic Research Science Publishers 369-372-bet 22-02-2025.
- 13.M Abdullayeva, Point spectrum of the operator matrices with the fredholm integral operators 2024/3/5, 47/47 Центр научных публикаций (buxdu. uz) 47 (47) BUXORO DAVLAT UNIVERSITETI ILMIY AXBOROTI
- 14.Р.Т.Мухитдинов, М.А.Абдуллаева КРАЙНИЕ ТОЧКИ МНОЖЕСТВА КВАДРАТИЧНЫХ ОПЕРАТОРОВ, ОПРЕДЕЛЕННЫХ НА S^1 "Scientific progress" 2021, 2/1, 470-477 ст.
Tog'ayev Turdimurod, Safarova Durдона, Hamidova Sevinch, Buvajonova Sevinch "Methods for solving odd-degree inverse equations." EXCELLENCIA: INTERNATIONAL MULTI-DISCIPLINARY JOURNAL OF EDUCATION <https://multijournals.org/index.php/excellencia-imje>